

Coburn

Em. Davis
Kelle. Cape

20
will

Boeing
Admiral

Al

Murphy

Tish
Jed

THE HETEROPHILE ANTIBODY

Patients acutely infected with EBV develop Paul-Bunnell-Davidsohn type heterophile antibodies (rabbit group). EB virus infections do not induce production of Forssman antibodies (guinea pig group). The differential heterophile absorption test for confirmation of the diagnosis of acute infectious mononucleosis is based on the fact that the heterophile antibody in infectious mononucleosis falls into the "rabbit" and not the "guinea pig" group. Of great interest is the fact that many Orientals fail to develop any heterophile antibodies what so ever when infected with EBV. Lymphoid cell lines obtained from Orientals, in contrast to Occidentals, usually express heterophile antigens which may explain why many Orientals do not develop heterophile antibodies.

SEROLOGIC REACTIONS OF TWO MAJOR CLASSES OF HETEROPHILE AB

<u>CLASS</u>	<u>DISEASE</u>	<u>REACTION WITH</u>			
		<u>HORSE</u> <u>RBC</u>	<u>SHEEP</u> <u>RBC</u>	<u>OX</u> <u>RBC</u>	<u>G. PIG</u> <u>KIDNEY</u>
Rabbit (P-B-D)	Mono	POS+	POS+	POS+	NEG-
Guinea Pig (Forssman)	Serum sickness	POS+	POS+	NEG-	POS+

WHEN SHOULD YOU ORDER EBV-SPECIFIC SEROLOGIC TESTS?

Order specific tests for IgG and IgM antibodies to EBV Capsid Antigen (Anti-VCA, IgG & Anti-VCA, IgM), and IgG antibodies to Nuclear Antigen (Anti-EBNA), in patients with negative heterophile tests who have a blood smear or a clinical picture suggestive of infectious mononucleosis. This is especially true in adults in whom typical clinical signs and symptoms are not always present, and in infants and children in whom typical heterophile responses may not occur. The heterophile test is not perfect and a more certain diagnosis can be made when specific EBV serologic tests are ordered. In addition, consider ordering serologic tests for CMV and Toxoplasmosis when the heterophile test is negative but the clinical findings suggest infectious mononucleosis.

SENSITIVITY AND SPECIFICITY OF HETEROPHILE TEST WITH DIFFERENTIAL ABSORPTION

<u>PATIENT GROUP</u>	<u>SENSITIVITY</u>	<u>SPECIFICITY</u>
Adolescents	96 %	93 %
Adults >40 years	89 %	93 %

IS THERE A SYNDROME ASSOCIATED WITH REACTIVATION OF EBV?

Because the virus resides in the infected host for life, it's always possible for the virus to begin replicating again. Renewed viral replication may be associated with recurrence of the disease. Today, specific serologic tests are available that usually respond only when the virus is replicating. Active viral replication is usually associated with a high titer of IgG antibody to the viral capsid antigen (>160) and antibody to early antigen. ^{1,2,5}

Exchano

Center S

Bent
NOC

Jo
Hatch

Garden

Johns for
Bleeds

Cleaner

Harold
S. Stevens
Applicant

Skipped
S. Stevens
Applicant

Double
Bill

Johns
for
Bleeds

Turners



ENCYCLOPAEDIA BRITANNICA • USA

OFFICE OF THE PRESIDENT

Dear Friend:

Recently we received a request from someone in your family for a free descriptive booklet and other information on The New Encyclopaedia Britannica. We want to thank you for your interest in this unique and exciting product, and make certain that you received the information you requested.

We also want to tell you about a rare opportunity through which you can obtain The New Encyclopaedia Britannica at a drastically reduced price.

Please read the reply card carefully, and answer the few brief questions. It will enable us to make sure that you received all the information and, at the same time, tell you of this unique offer.

You see, we sometimes have sets of Britannica that are no more than two years old, and are in excellent condition. Depending on availability, we can offer them at drastically reduced prices while the supply lasts.

Once again, we thank you for your interest. And if there is any other way we can be of service to you, please let us know.

Sincerely,

A handwritten signature in cursive script, reading 'Patricia A. Wier'.

Patricia A. Wier
President -EBUSA

1500

W. J. J. J.

Northway

Alen
admission
| Dusky
Service
Garage
Service
Station

W. J. J. J.

Houses
House

Bill
Sullivan
Daisy
Queen
House

Main St.

The syndrome of chronic mononucleosis is characterized by disabling fatigue and malaise, low-grade afternoon fever, and variable nonspecific symptoms including sore throat. The results of serologic tests include: a negative heterophile titer, and positive tests for Anti-VCA IgG, Anti-EA, and Anti-EBNA.

SEROLOGIC TEST RESULTS FOR EBV IN CHRONIC MONONUCLEOSIS

- Positive Anti-VCA, IgG, titer >160
- Positive Anti-EA, titer >5
- Positive Anti-EBNA
- Negative Anti-VCA, IgM
- Negative Heterophile Screening Test

SEROLOGIC TESTS FOR CHRONIC MONONUCLEOSIS ARE NOT SPECIFIC

Though there is little doubt about the occurrence of chronic EBV infections, there is doubt about the usefulness of routine serologic tests when making a diagnosis of chronic infectious mononucleosis.^{4,6} For example, elevated antibody titers (>40) to early antigen are found in as many as 13 percent of clinically healthy adults. Thus, the specificity of the test for antibodies to early antigen--when searching for chronic infectious mononucleosis-- is only 87 percent. Since clinically significant chronic infections with EBV are rare, but highly fatigued patients are common, most tired patients who test positive for antibodies to early antigen are false positives.^{6,7}

CELLULAR IMMUNITY AND EBV INFECTION

Our body's control of EBV is dependent upon a competent cellular immune system--not on humoral immunity. Thus, EBV serologic tests would not be expected to be highly sensitive or specific for chronic disease. In this regard, there is growing evidence that patients with chronic EBV infection may have a selective impairment of cellular immunity. Presumably, they have a defective recognition system for certain components of the nuclear antigen. These patients have extremely high titers to the Viral Capsid Antigen.¹

REFERENCES

1. Schooley RT, Carey RW, Miller G, Henle W, Eastman R, Mark EJ, Kenyon K, Wheeler EO, Rubin RH. Chronic Epstein-Barr virus infection associated with fever and interstitial pneumonitis. Clinical and serological features and response to antiviral chemotherapy. *Ann Intern Med* (1986)104:636-643.
2. Jones JF, Ray G, Minnich LL, Hicks MJ, Kibler R, Lucas DO. Evidence of active Epstein-Barr virus infection in patients with persistent unexplained illnesses: elevated anti-early antigen antibodies. *Ann Intern Med* (1985)102:1-7.
3. DuBois RE, Seeley JK, Brus I, Sakamoto K, Ballow M, Harada S, Bechtold TA, Pearson G, Purtilo DT. Chronic mononucleosis syndrome. *South Med J* (1984)77:1376-1382.
4. Merlin TL. Chronic mononucleosis: pitfalls in the laboratory diagnosis. *Human Pathol* (1986)17:1-8.
5. Straus SE, Tosato G, Armstrong G, Lawley T, Preble OT, Henle W, Davey R, Pearson G, Epstein J, Brus I, Blaese RM. Persisting illness and fatigue in adults with evidence of Epstein-Barr virus infection. *Ann Intern Med* (1985)102:7-16.
6. Tobi M, Straus SE. Chronic Epstein-Barr virus disease: a workshop held by the national institute of allergy and infectious diseases. *Ann Intern Med* (1985)103:951-953.
7. Greenberg DB. Depression, anxiety, and Epstein-Barr virus infection. *Ann Int Med* (1986)104:449.

M0367(7)



center st

hatch
home

Barn
~~garage~~
Cinder

Bill
Johnson
black

Washable Cleaner
Cle
Harold Stevens Radio Shop
OpStems
home

100 St

Ferry's
Note 199

Ken Durant

Winterrose
Mortuary

Main S



Eli Lilly and Company
Pharmaceutical Division

Lilly Corporate Center
Indianapolis, Indiana 46285
(317) 261-2000

December 1, 1986

Dear Doctor:

Eli Lilly and Company is pleased to announce the introduction of an important packaging innovation for its products in capsule dosage forms.

Capsules of Lilly/Dista products are now sealed by means of a unique banding process involving the application of a clear gelatin band that seals the halves of each capsule together.

The new process makes it difficult, if not impossible, to separate capsules without causing noticeable damage, thereby improving tamper-resistance. Although no dosage form, including tablets, can be made 100% tamperproof, Lilly believes its new capsule-banding process provides the consumer with a capsule that is as tamper-resistant as is possible to achieve.

Lilly/Dista products sealed with the new process are being phased in beginning with lots manufactured in September 1986.

As an innovative and responsible manufacturer of ethical pharmaceuticals, Lilly has always had as its primary goal the development of new and better products to meet the needs of patients. It is our hope that our most recent technologic advance in packaging will improve consumer safety and confidence in encapsulated medicines while enabling the proven advantages of this preferred dosage form to be retained.

Very truly yours,

PHARMACEUTICAL DIVISION
ELI LILLY AND COMPANY